


1. A 6.3 kg block sits on a table. A rope attached to the top of it exerts an upward force of 15.8 N on the box.

4

(a) _____ What is the weight W of the block?

- A)** 0.64 N **B)** 6.3 N **C)** 62 N



4

(b) _____ What is the normal force of the table on the block?

- A)** 15.8 N **B)** 45.9 N **C)** 61.7 N **D)** 77.8 N

4

2. _____ Which of the following statements is impossible (i.e. never true)?


- A)** "Ignoring air resistance, a ball thrown across the room is in free-fall."
B) "A car drives around a circle with constant velocity."
C) "A falling object has positive velocity."

4

3. _____ A ball is thrown into the air. At the top of its flight, the acceleration of the ball

- A)** points up **B)** is zero **C)** points down

4. A sailboat starts off at time $t = 0$ moving 6 m/s to the right, but a wind blows on the sailboat, giving it an acceleration of 2 m/s^2 to the left, eventually making the boat turn around. At time t , the boat is moving at 4 m/s to the left.



Δx	
v_i	
v_f	
a	
Δt	


(a) Fill in the table with what is given.

(b) What is the value of Δt ?

4

4

5. Two masses, 1 kg and 5 kg , are dropped from the same height. Ignore air resistance.



(a) _____ Which block feels a greater net force?

- A)** The 1 kg block **B)** The 5 kg block
- C)** Both feel the same nonzero force.
- D)** Both feel zero force.

(b) _____ Which block has the larger acceleration?

- A)** The 1 kg block **B)** The 5 kg block
- C)** Both have the same acceleration.

(c) _____ Which block will hit the ground first?

- A)** The 1 kg block **B)** The 5 kg block
- C)** Both will hit at the same time.

2

2


2

6. A ball is fired into the air. It reaches the top of its arc 10 m above the ground and 5 m along the ground. Suppose I want to know how fast the ball is moving at the top of the arc.

4

(a) Fill in everything that is known about the problem.

Δx		Δy	
v_{ix}		v_{iy}	
v_{fx}		v_{fy}	
a_x		a_y	
	Δt		



4


(b) _____ What is the first equation you would use to solve this equation?

- A) $v_{fx}^2 = v_{ix}^2 + 2a\Delta x$
- B) $\Delta y = \frac{1}{2}(v_{iy} + v_{fy})\Delta t$
- C) $v_{fx} = v_{ix} + a_x\Delta t$
- D) $\Delta y = v_{fy}\Delta t - \frac{1}{2}a_y(\Delta t)^2$
- E) $\Delta x = v_{ix}\Delta t + \frac{1}{2}a_x(\Delta t)^2$

4

7. _____ Two cannonballs are fired from different cannons at the same time. They both reach the same height, and hit different targets. Which target is hit first?


- A) A
- B) B
- C) Both at the same time



4

8. _____ A car is driving down the road at a constant 20 mph. The *net force* on the car
A) points forward **B)** is zero **C)** points backward

9. A person pushes horizontally on a 900 kg block with a force of 400 N. The block does not move. The coefficients of friction between box and floor are $\mu_s = 0.3$ and $\mu_k = 0.2$.



4

- (a) _____ What is the force of static friction on the block from the floor?

A) 180 N **B)** 270 N **C)** 400 N **D)** 1800 N **E)** 2600 N



4

- (b) _____ How hard would the person have to push to make the block move?


A) 180 N **B)** 270 N **C)** 400 N **D)** 1800 N **E)** 2600 N

4

10. A sign is pressed against a wall, attached by a rope. Find all the forces acting on the sign, and draw and label them on the rectangle below. Use the labels W , N , K , S , and/or T . I've given you one: the wall exerts an upward static frictional force on the sign.



11. A block is being pushed across the table by a finger, which exerts a normal force of $N_f = 8\text{ N}$ on the block. (The block's speed is not necessarily constant!) The table exerts a normal force upward of $N_t = 12\text{ N}$ on the block. Kinetic friction also acts on the block, with a coefficient of kinetic friction $\mu_K = 0.5$.



4


- (a) _____ What is the force of kinetic friction on the block?
A) 0.5 N **B)** 4 N **C)** 6 N **D)** 8 N **E)** 24 N

4

- (b) _____ This block is
A) speeding up **B)** slowing down
C) maintaining a constant speed

4



12. _____ Two blocks are stacked on the floor of an elevator. According to Newton's Third Law, Which of the following *must* have the same magnitude as the force of the floor on block Y?
- A) The force of block Y on the floor
 - B) The tension in the rope
 - C) The weight of block Y
 - D) The weight of both blocks
 - E) The force of block X on block Y



4

13. A block with mass 5 kg and weight 49 N is sliding down a frictionless hill with a $\theta = 50^\circ$ slope.

- (a) _____ Which is the correct force diagram for the block?



Box	x	y
$m\vec{a}$		


4

- (b) _____ Find the acceleration of the block.

- A) 0 m/s^2 B) 6.3 m/s^2
- C) 7.5 m/s^2 D) 9.8 m/s^2

4

14. _____ A block with a mass of 4 kg and a weight of 40 N is being lowered on a string. The block's acceleration is $a = 3 \text{ m/s}^2$ pointing upward. What is the tension in the rope?
- A) 28 N B) 40 N C) 52 N




15. The figure shows a stationary block of *weight* 90 N being pulled on by a rope which makes a 50° angle with the horizontal. The tension in the rope is 28 N.

4

(a) Fill in the table with what is given.

<i>Box</i>	<i>x</i>	<i>y</i>
Weight		
Normal, table		
Static Friction, table		
Tension, rope		




4

(b) Find the value of the static friction of the ground on the block.

16. A marble spins around the inside of a glass with a radius of 0.2 cm. The marble spins with a constant speed of 4 m/s.

4

(a) Calculate the linear acceleration a of the marble.



4

(b) _____ What type of force is acting on the marble to cause its circular motion?

A) normal **B)** tension **C)** static friction **D)** none of these